

Total RNA extraction from *Oryza sativa* leaf and *Solanum tuberosum* leaf using Ribospin™ Seed/Fruit

Experimental Conditions

Materials Required

- ◆ Ribospin™ Seed/Fruit (317-150, 50 preps)
- ◆ Tissuelyser II (85300, Supplier: Q) or another bead beating device
- ◆ Liquid nitrogen (LN₂)
- ◆ Absolute ethanol (C₂H₆O, CAS No. : 64-17-5, ≥99.0%)
- ◆ β-mercaptoethanol (C₂H₆OS CAS No. : 60-24-2, ≥99.0%)
- ◆ 1.5 ml or 2.0 ml microcentrifuge tube
- ◆ Vortex mixer
- ◆ Centrifuge (Max. speed 14,000 rpm or ≥10,000 x g)
- ◆ Pipette & sterile pipette tips
- ◆ Suitable protector (ex. lab coat, disposable gloves, goggles, etc.)
- ◆ Ice (for incubation)

Sample Information

Sample type :



Oryza sativa leaf



Solanum tuberosum leaf

- ◆ Sampling : After collecting fresh leaves, put them in a bag and seal it.
- ◆ How to store : Store in -70°C deep freezer
- ◆ Homogenizing : Tissuelyser II (85300, Supplier : Q)
- ◆ Conditions :
 - Sample amount :
 - Oryza sativa* leaf : 100 mg
 - Solanum tuberosum* leaf : 80 mg
 - Elution volume : 50 μl

Protocol

Before experiment

1. Before using for the first time, add absolute ethanol (ACS grade or better) into Buffer RBW and RNW as indicated on the bottle.
2. Prepare DNase I reaction mixture just before step 6.
 - Prepare aliquot DNase I and thaw on ice.
 - Mix 2 μl DNase I with 70 μl Buffer DRB.

Sample preparation

1. After measuring 100 mg of *Oryza sativa* leaf sample and 80 mg of *Solanum tuberosum* leaf sample, put them into a 1.5 ml microcentrifuge tube. Then, it is rapidly frozen using LN₂.
2. Grind the frozen sample using Tissuelyser II.
 - *Oryza sativa* leaf samples : 30 Hz, 40 sec
 - *Solanum tuberosum* leaf samples : 30 Hz, 20 sec
3. The next step is according to protocol II (not protocol I) of Ribospin™ Seed/Fruit.

Ribospin™ Seed/Fruit protocol II

* For more details, refer to the handbook of Ribospin™ Seed/Fruit

1. Add 500 μl Buffer SL and 5 μl β-mercaptoethanol to the sample and vortex vigorously for 15 sec.
2. Incubate the mixture for 3 min at room temperature and centrifuge the lysate at 13,000 rpm for 1 min. Transfer 300 μl of the supernatant to a new 1.5 ml microcentrifuge tube (not provided).
3. Add 300 μl Buffer ML to the supernatant and vortex vigorously for 15 sec and transfer all of the mixture to EzPure™ Filter (yellow).
4. Centrifuge at 13,000 rpm for 1 min and transfer 500 μl of the pass-through to a new 1.5 ml microcentrifuge tube (not provided).
5. Add 250 μl absolute ethanol to the supernatant and mix it well by inversion. Apply all of the mixture into Column Type F (blue ring) and centrifuge at 13,000 rpm for 1 min.
6. Add 500 μl Buffer RBW to Column Type F and centrifuge at 13,000 rpm for 30 sec.
7. Apply 70 μl DNase I reaction mixture onto the center of Column Type F for gDNA digestion. Incubate for 10 min at room temperature.
8. Add 500 μl Buffer RBW to Column Type F and centrifuge at 13,000 rpm for 30 sec. Add 500 μl Buffer RNW to Column Type F and centrifuge at 13,000 rpm for 30 sec.
9. Centrifuge at maximum speed for an additional 1 min to remove residual wash buffer. Transfer Column Type F to a new 1.5 ml microcentrifuge tube (provided).
10. Add 50 μl Nuclease-free water to the center of the membrane in Column Type F and centrifuge at 13,000 rpm for 1 min.

Total RNA extraction from *Oryza sativa* leaf and *Solanum tuberosum* leaf using Ribospin™ Seed/Fruit

Result

#	Sample	Conc. (ng/μl)	A _{260/280}	A _{260/230}	Yield (μg)
1	<i>Oryza sativa</i> leaf (100 mg)	183.4	2.13	2.35	9.17
2		173.4	2.14	2.38	8.67
3		164.3	2.15	2.35	8.22
4	<i>Solanum tuberosum</i> leaf (80 mg)	111.9	2.15	2.20	5.56
5		106.9	2.16	2.18	5.35

Table 1. The concentrations, yield and and purity of RNA extracted from 100 mg of *Oryza sativa* leaf and 80 mg of *Solanum tuberosum* leaf samples.

※ Remark
Absorbance measurement instrument : NanoDrop™ 2000/2000 c (ND-2000, Supplier : T)




Figure 1. The result of electrophoresis of total RNA from 100 mg of *Oryza sativa* leaf and 80 mg of *Solanum tuberosum* leaf samples.

Lane M : GENESTA™ 1 kb DNA Ladder with 5X loading dye (GA-100, GeneAll®, 1 μl loading)

Lanes 1~3 : 100 mg of *Oryza sativa* leaf samples (3 μl loading)

Lanes 4~5 : 80 mg of *Solanum tuberosum* leaf samples (5 μl loading)

※ Remark
Electrophoresis conditions : 1.0% agarose gel (150 V, 17 min)