

AmpONE™ 2X Taq Premix

Ver 3.1

Cat. No. 526-200 (20 µl PCR reaction)

Storage at -20°C

Disclaimer

For research use only. Not for use in diagnostic or therapeutic procedures.

Description

AmpONE™ 2X Taq Premix is made from GeneAll® AmpONE™ Taq DNA Polymerase which is purified from the cloned *Thermus aquaticus* DNA polymerase gene in *Escherichia coli* (*E. coli*). This premix contains all reaction components required for conventional PCR, such as reaction buffer, dNTP, loading dye, stabilizer and sediment in addition to Taq DNA polymerase. It is recommended for use in conventional PCR (below 5 kb), TA cloning and colony PCR.

AmpONE™ 2X Taq Premix is highly processive 5' → 3' DNA polymerase that lacks 3' → 5' exonuclease (proofreading) activity. This premix is pre-aliquoted in 8-strip tube and is stable for 18 months at -20°C. Therefore, this premix serves time-saving, cost-effective experiment.

Quality Control

No endonuclease activity, nicking activity, exonuclease activity, or priming activity has been detected.

Components

Cat. No.	Reaction Vol.	AmpONE™ 2X Taq Premix *
526-200	20 µl	8-strip tube x 12 ea with tube rack

* AmpONE™ 2X Taq Premix contains loading dye.

Storage Conditions

Stable for 18 months when all components are stored in a frost-free freezer at -20°C.

Features

High efficiency, ready-to-use, time-saving, pre-aliquot type, and cost-effective

Applications

Conventional PCR (up to 5 kb), TA cloning, Colony PCR, High through-put PCR, Routine PCR requiring high reproducibility, DNA sequencing template preparation

Note

Do not contaminate the AmpONE™ 2X Taq Premix with primers and template DNA used in individual reactions. Thaw and mix all components thoroughly, spin down shortly and chill on ice.

Recommended PCR Mixture

The volume of components for 20 µl PCR reaction.

Components	Reaction Volume
	20 µl
AmpONE™ 2X Taq Premix	10 µl
Forward primer (10 pmole/µl)	1 µl
Reverse primer (10 pmole/µl)	1 µl
Template DNA *	-
Add D.W. to	20 µl

* Recommended amount of template per PCR reaction:

- < 50 ng plasmid or
- < 500~1,000 ng genomic DNA or
- < 2 µl of a 100 µl single plaque eluate or
- 1 single bacterial colony

PCR Condition

Step	Temp.	Time	Cycles
Initial Denaturation	95°C	5 min	1
Denaturation	95°C	30 sec	25~40
Annealing	A°C	30 sec	
Extension	72°C	B	
Final Extension	72°C	5 min	1
Cooling	4°C	∞	-

A: Recommended annealing temperature: 50~65°C

The value is 4-6 lower than T_m of primers

$$T_m = 2 (A+T) + 4 (G+C)$$

B: Recommended extension time: 10~60 sec

1 min/kb (the size of target PCR product)